

Quantum Transport Evidence for the Three-Dimensional Dirac Semimetal Phase in Cd_3As_2 and Observation of Superconductivity under High Pressure

L. P. He¹, Y. T. Jia², S. J. Zhang², X. C. Hong¹, J. K. Dong¹, J. Pan¹, Z. Zhang¹, J. Zhang¹, C. Q. Jin^{2*} and S. Y. Li^{1**}

¹Department of Physics, Surface Physics Laboratory (National Key Laboratory), and Laboratory of Advanced Materials, Fudan University, Shanghai 200433, China


²Beijing National Laboratory for Condensed Matter Physics and Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China




Abstract

We report the quantum transport properties of Cd_3As_2 single crystals in a magnetic field. A large linear quantum magnetoresistance is observed near room temperature. With decreasing temperature, the Shubnikov—de Haas oscillations appear in both the longitudinal resistance R_{xx} and the transverse Hall resistance R_{xy} . From the strong oscillatory component ΔR_{xx} , a linear dependence of the Landau index n on $1/B$ is obtained, and it gives an n -axis intercept between $1/2$ and $5/8$. This clearly reveals a nontrivial π Berry's phase, which is a distinguished feature of Dirac fermions. And the resistance of Cd_3As_2 under pressure up to 50.9 GPa was also measured. Surprisingly, superconductivity with $T_c \approx 2.0$ K emerges at 8.5 GPa. The T_c keeps increasing to about 4.0 K at 21.3 GPa, then shows an anomalous nearly constant pressure dependence up to the highest pressure 50.9 GPa. Our observation of superconductivity in pressurized three-dimensional Dirac semimetal Cd_3As_2 provides an interesting candidate for topological superconductor.


Magnetoresistance




Berry's Phase



Superconductivity



H_{c2} and Phase Diagram



References

- [1] Z. K. Liu, J. Juan *et al.*, *Nat. Mater.* **13**, 677 (2014).
- [2] Y. B. Zhang, Y.W. Tan, H. L. Stormer, and P. Kim, *Nature (London)* **438**, 201 (2005).
- [3] H. Murakawa *et al.*, *Science* **342**, 1490 (2013).
- [4] K. Kirshenbaum *et al.*, *Phys. Rev. Lett.* **111**, 087001 (2013).

Summary

In summary, we have performed bulk transport measurements on single crystals of the proposed 3D Dirac semimetal Cd_3As_2 . A large linear quantum magnetoresistance is observed near-room temperature. By analyzing the Shubnikov—de Haas oscillations of longitudinal resistance at low temperature, a nontrivial π Berry's phase with a small phase shift is obtained, which provides bulk quantum transport evidence for the existence of a 3D Dirac semimetal phase in Cd_3As_2 . We have done resistance measurements on the 3D Dirac semimetal Cd_3As_2 single crystals under pressures up to 50.9 GPa. Below 6.4 GPa, the resistance behavior becomes more and more insulating with increasing pressure, however it changes back to metallic again at higher pressures. Superconductivity emerges at 8.5 GPa. The T_c increases from 2.0 K at 8.5 GPa to 4.0 K at 21.3 GPa, then it shows an anomalous constant pressure dependence up to the highest pressure measured.