

# Geometric control of optical dipoles at a cold atom--nanofiber interface



Ruijuan Liu<sup>1,\*</sup>, Jinggu Wu<sup>1</sup>, Yudi Ma<sup>1</sup>, Yanting Zhao<sup>2,†</sup>, Saijun Wu<sup>1,§</sup>,


<sup>1</sup>State Key Laboratory of Surface Physics and Department of Physics, Fudan University, Shanghai, People's Republic of China

<sup>2</sup>State Key Laboratory of Quantum Optics and Quantum Optics Devices, Institute of Laser Spectroscopy, Shanxi University, Taiyuan, 030006, China

## Introduction:


- Novel quantum optics research requires precise, arbitrary control of optical dipoles.
- Nanophotonic interfaces: Strong interaction between confined photons and atoms.
- Challenge: How to perfect optical control in the near field, despite the field inhomogeneity?
- This work: We demonstrate geometric phase control of optical dipoles at a nanoscale atom-fiber interface with a high efficiency agreeing with theoretical prediction. The robust technique can be perfected to support near-field-lattice based 1D quantum optical researches.

### Evanescent coupling and control at the cold atom-nanofiber interface



### Geometric phase control of optical dipoles



- Electric dipole moment for 2-level atom:  $\langle \mathbf{d} \rangle = \mathbf{d}_{eg} \rho_{ge} + c.c. \Rightarrow \mathbf{d}_{eg} \rho_{ge} \boldsymbol{\eta}_d e^{-i\gamma} + c.c.$
- The geometric phase  $\gamma$  can be written to the ground state  $|g\rangle$  by cyclically driving the auxiliary  $|a\rangle \rightarrow |a\rangle$  transition.  $\boldsymbol{\eta}_d$ : dipole control complex coefficient.




## Measurement Principles:

### Evanescent attenuation of guided probe by cold atoms

**Optical theorem:** The scattering of the guided probe by the atoms is proportional to the imaginary part of the coherent forward scattering into the same mode.




### What happens if the relative phase between the dipole and probe suddenly jumps?





## Experimental Setup:

### Cold atom + ONF interface with 2D+MOT loading




### Probe and control waveforms




## Robust population inversion:

### Inversion by Rapid Adiabatic Passage [1,2]





- Efficient D1 population inversion can be driven by either composite[2] or adiabatic[1] pulses within a nanosecond at ONF interface.
- The experimental observations agree excellently with full-level numerical simulation in the near field.

## Geometric control of optical dipole:




- By successively applying two optimally chirped pulses with  $\phi_{21}$  phase difference, the optical phase is transferred to the near-field atomic dipoles,  $\gamma = \phi_{12} + \pi$ , reflected in the transient probe transmission.
- A probe phase-jump globally induces the same  $\phi_{21}$  relative to  $\mathbf{E}_s^f$ , leading to similar transmission signal.
- According to  $\mathbf{E}_s(\mathbf{r}) \sim \sum \mathbf{G}(\mathbf{r}, \mathbf{r}_j) \cdot \mathbf{d}_j$ , we estimate the dipole control efficiency by observing the  $\mathbf{E}_s$ -transient within nanoseconds.

### Experimental results



### Numerical model



- Keep it simple: uniform, non-interacting atoms in the near field.
- HE11 mode, evanescent full-level OBE.
- Parameters match the experiment.

- Exp geometric phase control via evanescent couplings with an efficiency averaged over the near field dipoles to be above 50%.
- Exp performance is about 20% lower than numerical expectation, likely related to inaccurate near-field modeling of distributions.
- Numerical model predicts atomic dipole control with 90% fidelity at specific locations with linear local polarization and mF=0.

## Summary & outlook

- We demonstrate geometric phase control of optical dipoles at a cold atom-nanofiber interface, for the first time. Ensemble-averaged efficiency  $|\eta_d| \approx 50\%$  is retrieved with a phase-jump spectroscopy. The result is agreeable with theoretical expectations for the nanoscale quantum control.
- Novel 2DMOT-ONF interface with ~field-free line.
- In our next step, by loading atoms into a near-field lattice and by sending geometric-phase-writing ONF-guided pulses from opposite directions, the D2 collective dipole excitation can be coherently shifted into sub-radiant domain, potentially enabling highly nontrivial many-body quantum optical researches.

## Reference

- Y. He et al, "Geometric Control of Collective Spontaneous Emission", *Phys. Rev. Lett.* 125, 213602 (2020)
- Y. Ma et al, "Composite picosecond control of atomic state through a nanofiber interface", *Phys. Rev. Applied* 20, 024041 (2023)